




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C01B 3/00, C11D 3/39, 7/38, C07D 265/12, 295/12, 295/14, 211/26, 211/34		A1	(11) International Publication Number: WO 98/23533
			(43) International Publication Date: 4 June 1998 (04.06.98)
(21) International Application Number: PCT/US97/21335		(81) Designated States: AU, BR, CA, CN, ID, JP, KR, MX, PL, RU, SG, TR, UA, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 18 November 1997 (18.11.97)			
(30) Priority Data: 08/758,540 29 November 1996 (29.11.96) US		Published <i>With international search report.</i>	
(71) Applicant: THE CLOROX COMPANY [US/US]; 1221 Broadway, Oakland, CA 94612 (US).			
(72) Inventors: CHOY, Clement, K.; 1345 Sugarloaf Drive, Alamo, CA 94507 (US). DELINE, James, E.; 3857 MacGregor Common, Livermore, CA 94550 (US). FOLAND, Lafayette, D.; 11784 Bloomington Way, Dublin, CA 94568 (US). GARABEDIAN, Aram, Jr.; 37805 Elliott Street, Fremont, CA 94536 (US). KLOTTER, Kevin, A.; 1349 Cameo Court, Livermore, CA 94550 (US). PETRIN, Michael, J.; 3613 Old Glory Court, Antioch, CA 94508 (US). PHILLIPPI, Martin, A.; 3641 Dunsmuir Circle, Pleasanton, CA 94588 (US). SMITH, William, L.; 3227 Runnymede Court, Pleasanton, CA 94588 (US).			
(74) Agents: HAYASHIDA, Joel, J. et al.; 1221 Broadway, Oakland, CA 94612 (US).			

(54) Title: **LIQUID COMPOSITIONS CONTAINING N-ALKYL AMMONIUM ACETONITRILE SALTS**

(57) Abstract

The invention provides liquid cleaning or bleaching compositions containing compounds having the structure of Formula (I) wherein A is a saturated ring formed by five atoms in addition to the N₁ atom, the five saturated ring atoms being four carbon atoms and a heteroatom, the substituent



R₁ bound to the N₁ atom of the Formula (I) structure including either (a) a C₁₋₂₄ alkyl or alkoxyalkyl where the alkoxy is C₂₋₄, (b) a C₄₋₂₄ cycloalkyl, (c) a C₇₋₂₄ alkaryl, (d) a repeating or nonrepeating alkoxy or alkoxyalkyl alcohol, where the alkoxy unit is C₂₋₄, or (e) —CR₂R₃CN where R₂ and R₃ are each H, a C₁₋₂₄ alkyl, cycloalkyl, or alkaryl, or a repeating or nonrepeating alkoxy or alkoxyalkyl alcohol where the alkoxy unit is C₂₋₄, the R₂ and R₃ substituents are each H, a C₁₋₂₄ alkyl, cycloalkyl, or alkaryl, or a repeating or nonrepeating alkoxy or alkoxyalkyl alcohol where the alkoxy unit is C₂₋₄, and wherein Y is a generally anionic counterion and Z is in the range of 0 to 10. Additional, adjunct materials commonly found in liquid cleaning and bleaching compositions can be added.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

LIQUID COMPOSITIONS CONTAINING N-ALKYL
AMMONIUM ACETONITRILE SALTS

5

Field of the Invention

10 The present invention generally relates to N-alkyl ammonium acetonitrile compounds in a liquid matrix, and particularly liquid compositions containing a source of active oxygen useful in applications such as bleaching and cleaning.

 This application is a continuation in part of Serial No. 08/475,292, filed June 7, 1995, entitled "N-ALKYL AMMONIUM ACETONITRILE BLEACH ACTIVATORS," inventors Arbogast et al., of common assignment herewith.

15

Background of the Invention

 Peroxy compounds are effective bleaching agents, and compositions including mono- or di-peroxyacid compounds are useful for industrial or home cleaning or laundering operations. For example, U.S. Patent 3,996,152, issued December 7, 1976, inventors Edwards et al., discloses bleaching compositions including peroxygen compounds such as diperazelaic acid and diperisophthalic acid.

20

 Peroxyacids (also known as "peracids") have typically been prepared by the reaction of carboxylic acids with hydrogen peroxide in the presence of sulfuric acid. For example, U.S. Patent 4,337,213, inventors Marynowski et al., issued June 29, 1982, discloses a method for making diperoxyacids in which a high solids throughput may be achieved.

25

 However, bleaching products containing peroxyacid compounds tend to lose bleaching activity during storage, due to decomposition of the peroxyacid. The relative instability of peroxyacid can present a problem of storage stability for compositions consisting of or including peroxyacids.

2

One approach to the problem of reduced bleaching activity of peroxyacid compositions has been to include activators of peroxyacids. U.S. Patent 4,772,290, issued September 20, 1988, inventors Mitchell et al., and U.S. Patent 4,900,469, issued February 13, 1990, inventors Farr et al., both of which are of common assignment herewith, disclose stable organic bleach activators suspended in a liquid hydrogen peroxide matrix.

U.S. Patent 4,915,863, issued April 10, 1990, inventors Aoyagi et al., discloses compounds said to be peracid precursors that have nitrile moieties. U.S. Patent 5,236,616, issued August 17, 1993, inventors Oakes et al., discloses compounds said to be cationic peroxyacid precursors that have nitrile moieties. These nitrile containing activators do not contain a leaving group, such as found in ester-type bleach activators, but instead include a quaternary ammonium group suggested as activating the nitrile and said, upon hydrolysis in the presence of hydrogen peroxide, to generate a peroxyimide acid as bleaching species. The Aoyagi et al. activators include an aromatic ring, which tends to cause fabric yellowing.


German patent application DE OS 44 312 12, published March 7, 1996, describes production of quaternized glycinonitriles in the form of stable aqueous solutions.

In the parent application, Serial No. 08/475,292, N-methylmorpholinium acetonitrile methylsulfate ("MMA") is noted as being of special interest. The MMA is useful as an activator in bleaching applications when combined with a source of active oxygen. These compounds and related ones, however, may be incorporated into liquid formulations, both with and without a source of active oxygen.

Summary of the Invention

The present invention provides liquid cleaning or bleaching compositions comprising compounds having the structure of Formula I

FORMULA I




wherein A is a saturated ring formed by five atoms in addition to the N₁ atom, the five saturated ring atoms being four carbon atoms and a heteroatom, the substituent R₁ bound to the N₁ atom of the Formula I structure including either (a) a C₁₋₂₄ alkyl or alkoxyalkyl where the alkoxy is C₂₋₄, (b) a C₄₋₂₄ cycloalkyl, (c) a C₇₋₂₄ alkaryl, (d) a repeating or nonrepeating alkoxy or alkoxyalkyl alcohol, where the alkoxy unit is C₂₋₄, or (e) -CR₂R₃C≡N where R₂ and R₃ are each H, a C₁₋₂₄ alkyl, cycloalkyl, or alkaryl, or a repeating or nonrepeating alkoxy or alkoxyalkyl alcohol where the alkoxy unit is C₂₋₄, the R₂ and R₃ substituents are each H, a C₁₋₂₄ alkyl, cycloalkyl, or alkaryl, or a repeating or nonrepeating alkoxy or alkoxyalkyl alcohol where the alkoxy unit is C₂₋₄, and wherein Y is a generally anionic counterion and Z is in the range of 0 to 10.

The Formula I compounds have a quaternary nitrogen atom (N₁), requiring the presence of at least one counterion (Y) to be associated therewith. Counterions for the inventive salts include both organic and inorganic anions, preferably alkylsulfate, especially methylsulfate, sulfate or bisulfate, tosylate and mesylate, and chloride, bromide, and nitrate. Especially preferred are methylsulfate, sulfate, bisulfate, tosylate and mixtures thereof, which have been found to result in surprisingly advantageous properties for the novel Formula I compounds. Additionally, although Y has been indicated in Formula I to be monovalent, in fact, it can be multivalent. As an example, when Y is sulfate, the counterion is divalent. Accordingly, these inventive salts are particularly well suited to bleaching and cleaning compositions.

The novel compounds with the Formula I structure are particularly useful when formulated as compositions that include a source of active oxygen, and these compositions provide excellent bleaching in alkaline solutions.

Preferred embodiments of the invention include lower alkyls substituted at the N₁, e.g. N-methyl morpholinium acetonitrile, N-ethyl morpholinium acetonitrile, N-butyl morpholinium acetonitrile, which are illustrated by Formula II (with "n" preferably being 0 to 24 and where "Y" is one of the above described counterions).

FORMULA II



A particularly preferred embodiment of the invention are liquid compositions containing N-methyl morpholinium acetonitrile methylsulfate, sulfate, bisulfate salts (sometimes designated as, respectively, "MMAMS," "MMAS" or "MMABS," where "n" of Formula II is 0), or mixtures thereof, which have excellent stability, and which show excellent bleaching and cleaning performance when formulated with a source of active oxygen in alkaline wash water. Tosylate salts are also preferred and would be designated as "MMATS."

When formulated with a source of active oxygen, compositions of the invention are useful as or in home cleaning and laundry products, such as bleaching additives, detergents, detergent boosters, detergents with bleach, bleaches, bleaching aids, dishwashing detergents, surface and mildew stain removers, and spot treatment products such as stain removers, prewash and presoak laundry aids. Among the advantages derived from compositions of the invention are improved cleaning, stain removal, spot removal, whitening, and brightening of treated articles.


Brief Description of the Drawing

Fig. 1 shows a graph depicting performance of a preferred embodiment of the invention formulated as a hard surface cleaner at various pH's.

Detailed Description of the Preferred Embodiments

The novel compounds used in the invention include certain nitriles having the structure illustrated by Formula I. The N₁ atom of the Formula I compound is part of a saturated ring, illustrated by "A" in Formula I.


FORMULA I




This saturated ring of which N₁ is a part has five atoms in addition to N₁, with at least one heteroatom being in the saturated ring in addition to the N₁, preferably wherein the heteroatom of the ring is an oxygen atom or a sulfur atom.

The N₁ atom shown in Formula I is N-acetonitrile substituted and also quaternized. Without being bound by theory, it is believed that the electron withdrawing nature of the quaternary nitrogen may be increased by being part of a saturated, heterocyclic ring and may also function to improve the hydrophilic character of the oxidant.

A substituent R₁ will be bonded to the N₁ atom of the Formula I structure and additionally a nitrile moiety (-CR₂R₃C≡N) is bonded to the N₁ atom, where R₂ and R₃ are each H, a C₁₋₂₄ alkyl, cycloalkyl, or alkylaryl, or a repeating or nonrepeating alkoxy or alkoxyated alcohol where the alkoxy unit is C₂₋₄. The R₁ substituent may be a C₁₋₂₄ alkyl or alkoxyated alkyl where the alkoxy is C₂₋₄, a C₄₋₂₄ cycloalkyl, a C₇₋₂₄ alkylaryl, a repeating or nonrepeating alkoxy or alkoxyated alcohol, where the alkoxy unit is C₂₋₄, and illustrative such groups are, for example,




where j=1 to 24. The R₁ substituent may also be another --CR₂R₃C≡N, and again R₂ and R₃ are each H, a C₁₋₂₄ alkyl, cycloalkyl, or alkylaryl, or a repeating or nonrepeating alkoxy or alkoxyated alcohol where the alkoxy unit is C₂₋₄, and illustrative such groups are:



where j=1 to 24.

Particularly preferred activator embodiments are illustrated by Formula II (where "Y" is as earlier described and "n" is 0 to 24).

FORMULA II




Novel derivatives used in the liquid compositions of the invention include peroxyimidic intermediates that are believed formed from the novel nitriles in the presence of an active oxygen source. So formed, peroxyimidic derivatives typically

would be short-lived intermediates formed ⁶in situ when the nitriles of the invention interact with a source of hydrogen peroxide and where the reactive nitrile moiety forms a peroxyimidic acid. However, such peroxyimidic derivatives may also be prepared *in situ* by analogy to syntheses known in the art.

1. Counterions

Since compounds of the invention are typically quaternized, they will include at least one counterion (designated as "Y"), selected from a wide variety of anionic counterions, especially alkylsulfate (e.g., methylsulfate), sulfate or bisulfate, or mixtures thereof, mesylate and tosylate. In the copending parent application of which this is a continuation-in-part, N-alkyl ammonium acetonitrile compounds are disclosed as typically including such wide variety of counterions such as chloride, bromide, nitrate, alkyl sulfate, and the like. There, the preferred embodiment was described as N-methyl ammonium acetonitrile methyl sulfate. In addition, in the present invention of sulfate or bisulfate salt may be produced from heated and acidified MMA (as methyl sulfate). However, the sulfate and bisulfate salts appear to exist in an equilibrium, the predominance of one over the other dependent upon pH. Three preferred salts are illustrated by Formulas IIIA, IIIB and IIIC.

FORMULA III



2. Bleaching and Cleaning Compositions

Bleaching and cleaning compositions of the invention include the Formula I nitrile salts as activator, together with a source of active oxygen.

The peroxide or active oxygen source for compositions of the invention may be selected from most preferably hydrogen peroxide, Caro's acid (peroxymonosulfuric acid), and then, as suspended particulate oxidants, the alkali metal and alkaline metal salts of percarbonate, perborate, persilicate and hydrogen peroxide adducts. Examples of hydrogen peroxide formulations suitable for use herein include those depicted in Mitchell et al., U.S. Patent 4,900,468, Farr et al., U.S. Patent 5,180,514 and Baker et al., U.S. Patent 4,764,302, all of common assignment and all of which are incorporated herein by reference. Where sodium percarbonate, sodium perborate mono- and tetrahydrate, are utilized in aqueous formulations, it is most preferable to suspend them in such aqueous formulations, along with stabilizers. Exemplary of these systems are Peterson et al., U.S. Patent 5,464,552, Published European Patent Applications EP 294 904 and EP 293 040, incorporated herein by reference. Other peroxygen sources may be possible, such as monopersulfates and monoperphosphates, or their equivalent aqueous forms, such as monopersulfuric acid, known in the trade as Caro's acid or Caroate®, a product of BASF AG, Germany; and poorly soluble oxidants, such as alkaline earth peroxides, for example, Gray et al., U.S. Patents 4,891,147 and 5,019,189, both of which are incorporated herein by reference. Alternatively, the formulations of the invention may be essentially nonaqueous. These nonaqueous formulations will have a nonaqueous liquid as the continuous phase, such as nonionic surfactant, or nonaqueous organic solvents such as glycol ethers, hydrocarbons, acids, alcohols, and the like. A nonaqueous formulation would be expected to be highly chemically stable because bleach activators such as MMA are less likely to be hydrolyzed in the nonaqueous continuous matrix. Exemplary nonaqueous formulations are depicted in Peterson et al., U.S. Patent 4,874,537, and Van Buskirk et al., U.S. 5,415,796, both of which are incorporated herein by reference.

The range of peroxide to activator is preferably determined as a molar ratio of peroxide to activator. Thus, the range of peroxide to each activator is a molar ratio of from about 0.1:1 to 100:1, more preferably about 1:1 to 10:1 and most preferably about 2:1 to 8:1. This peracid activator/peroxide composition should provide about 0.5 to 100 ppm A.O., more preferably about 1 to 50 ppm peracid A.O. (active oxygen), and most preferably about 1 to 20 ppm peracid A.O., in aqueous media for typical laundry

8

applications. Formulations intended for hard surface cleaning will more typically have peracid activator/peroxide providing from about 0.5 to 500,000 ppm A.O., more preferably about 1 to 20,000 ppm peracid A.O., and most preferably about 1 to 10,000 ppm peracid A.O.

5 However, one of the advantages of the inventive MMA compounds is that those tested have been found to be oxidatively stable when formulated with hydrogen peroxide, preferably at a low pH. This allows for great flexibility when formulating liquid products since the MMA compounds do not necessarily need to be kept separate from the source of active oxygen.

10 Compositions of the invention have been found to provide superior bleaching (cleaning and stain removal) benefits on common laundry and household stains.

3. Delivery Systems

15 The liquid compositions of the invention can take numerous desirable forms. For example, without limitation, these include macroemulsions, microemulsions, structured liquids, liquid crystals, vesicular systems, lamellar systems, suspensions, dispersions, gels, mulls and pastes. These liquid systems can be clear or translucent (such as microemulsions) to opaque. These liquid systems can either be cleaning liquids wherein the inventive compounds are present as potentially antimicrobial actives, or, where a source of peroxygen (active oxygen) is present, as a bleaching liquid, as well as
20 detergents and detergent bleaches. They can also be either unitary systems, or multiple deliveries, such as, for example, a dual chamber container, one chamber containing the inventive N-alkyl ammonium acetonitrile compounds with a source of active oxygen, while the other contains a solution including actives sensitive to oxidation, for example, enzymes and fluorescent whitening agents. An example of a container which can
25 co-dispense these two different liquid compositions is found in Beacham et al., U.S. Patent 4,585,150, incorporated herein by reference; and an example of a system where a dual delivery is depicted, with one part containing a liquid oxidant formulation, the other, a liquid with materials which are sensitive to oxidation, is found in Co-pending applications of Choy et al., Serial Nos. 08/605,822, and 08/605,824, both filed February
30 23, 1996, and both entitled "Composition and Apparatus for Surface Cleaning," of common assignment, incorporated herein by reference. In the case of more viscous

9

forms, such as gels, mulls or pastes, the continuous phase can be nonionic surfactants, and is exemplified by Kaufmann et al., U.S. Patents 4,743,394 and 5,362,413, both of which are incorporated herein by reference.

5 The liquid compositions of this invention frequently contain varying amounts of surfactants, which may act both as a cleaning active ingredient, but also to help disperse sparingly soluble materials in the liquid phase, or which may serve as the continuous phase in an essentially nonaqueous composition.

4. Surfactants

10 Surfactants with which the activators and active oxygen compositions may be combined or admixed include linear ethoxylated alcohols, such as those sold by Shell Chemical Company under the brand name Neodol. Other suitable nonionic surfactants can include other linear ethoxylated alcohols with an average length of 6 to 16 carbon atoms and averaging about 2 to 20 moles of ethylene oxide per mole of alcohol; linear and
15 branched, primary and secondary ethoxylated, propoxylated alcohols with an average length of about 6 to 16 carbon atoms and averaging 0-10 moles of ethylene oxide and about 1 to 10 moles of propylene oxide per mole of alcohol; linear and branched alkylphenoxy (polyethoxy) alcohols, otherwise known as ethoxylated alkylphenols, with an average chain length of 8 to 16 carbon atoms and averaging 1.5 to 30 moles of ethylene oxide per mole of alcohol; and mixtures thereof. Shell Chemical, Huntsman
20 Chemical and Union Carbide are among the numerous producers of these surfactants.

Further suitable nonionic surfactants may include polyoxyethylene carboxylic acid esters, fatty acid glycerol esters, fatty acid and ethoxylated fatty acid alkanolamides, certain block copolymers of propylene oxide and ethylene oxide, and block polymers or propylene oxide and ethylene oxide with propoxylated ethylene diamine. Also included
25 are such semi-polar nonionic surfactants like amine oxides (such as Ammonyx from Stepan and Barlox from Lonza), phosphine oxides, sulfoxides and their ethoxylated derivatives.

Anionic surfactants may also be suitable. Examples of such anionic surfactants may include the ammonium, substituted ammonium (e.g., mono-di-, and triethanolammonium), alkali metal and alkaline earth metal salts of C₆-C₂₀ fatty acids and
30 rosin acids, linear and branched alkyl benzene sulfonates, alkyl sulfates, alkyl ether

10
sulfates, alkane sulfonates, alpha olefin sulfonates, hydroxyalkane sulfonates, fatty acid monoglyceride sulfates, alkyl glyceryl ether sulfates, acyl sarcosinates and acyl N-methyltaurides.

5
Suitable cationic surfactants may include the quaternary ammonium compounds in which typically one of the groups linked to the nitrogen atom is a C₁₂-C₁₈ alkyl group and the other three groups are short chained alkyl groups which may bear inert substituents such as phenyl groups.

10
Suitable amphoteric and zwitterionic surfactants containing an anionic water-solubilizing group, a cationic group or a hydrophobic organic group include amino carboxylic acids and their salts, amino dicarboxylic acids and their salts, alkyl-betaines, alkyl aminopropylbetaines, sulfobetaines, alkyl imidazolinium derivatives, certain quaternary ammonium compounds, certain quaternary phosphonium compounds and certain tertiary sulfonium compounds.

15
These and other types of surfactants are exemplified in McCutcheon's Emulsifiers and Detergents (1994) and Kirk-Othmer Encyclopedia of Chemical Technology 3rd, Vol.22, "Surfactants," pp. 332-432 (1983), both of which are incorporated herein by reference.

Other common detergent adjuncts may be added if a bleach or detergent bleach product is desired; See 6 below.

20 5. Source of Acid/Alkali

25
30
Compositions of the invention, when combined with a source of active oxygen, preferably function for bleaching best at a neutral to alkaline pH, but are shelf stabilized best at an acidic pH. Thus, compositions of the invention preferably include a source of protons as an "acid sink." This can be achieved by having a mineral or organic acid present. These include, but are not limited to, mineral acids selected from phosphoric, sulfuric, hydrochloric, nitric, carbonic, boric, sulfamic, sulfurous acids and mixtures thereof; and organic acids selected from acetic, hydroxyacetic (glycolic), citric, succinic, maleic benzoic, oxalic acids, and mixtures thereof. It may also be possible to use the acidic forms of materials ordinarily viewed as hydrotropes, such as toluene sulfonic acid, xylene sulfonic acid, cumene sulfonic acid, and the like; and surfactants, such as

alkylbenzene sulfonic acid (also known as HLAS), exemplary of which is dodecyl
benzene sulfonic acid (See Choy et al., U.S. Patents 4,759,867, 4,804,491 and 4,895,669,
incorporated herein by reference). However, the compositions of the invention can also
be pH adjusted for a variety of applications, such as when used as a mildew or other type
of stain remover.

When the composition is ready for use as a laundry additive, it is especially
advantageous to have an amount of alkaline buffer present sufficient to maintain a pH
greater than about 6.0, more preferably in the range of about 8.5 to about 10.5 for most
effective bleaching, when the liquid formulation is dispensed into an aqueous wash
system. As a hard surface cleaner, on the other hand, it may be useful to co-dispense the
alkaline buffer in a separate, preferably liquid, composition. These alkaline buffers
include, but are not limited to, alkali metal hydroxides (sodium, lithium, potassium),
ammonium hydroxide, alkali metal ortho-, meta- and pyrophosphates, alkali metal
silicates, alkali metal tetraborates (penta- and decahydrates), alkali metal and ammonium
carbonates, alkali metal and ammonium carbamates (See Garabedian, Jr., et al., U.S.
Patents 5,523,024, 5,468,423, 5,437,807 and 5,252,245, all incorporated herein by
reference), alkali metal and ammonium polyacrylates, alkali metal and ammonium
succinates, alkali metal and ammonium maleates and additional conjugate bases of weak
organic acids, such as those mentioned hereinabove. Further, organic bases are included,
such as, without limitation, ethanolamine, diethanolamine, triethanolamine,
hydroxyamine, methylamine, dimethylamine and trimethylamine. On the other hand,
acidic hard surface cleaners are certainly well known and preferred for use as bathroom
cleaners. These types of cleaners therefore are acidified with the type of acids described
hereinabove.

6. Additional Functional/Aesthetic Adjuncts

Other adjuncts (useful in cleaning and laundering applications) are optionally
included in the inventive compositions. Dyes include anthraquinone and similar blue
dyes. Pigments may also be used. Monastral colorants are also possible for inclusion.
Brighteners or whiteners, such as stilbene, styrene and styrylnaphthalene brighteners
(fluorescent whitening agents), may be included. Fragrances used for aesthetic purposes
are commercially available from Quest, Sozio, Firmenich, Dragoco, Bush Broke and
Allen, Norda, International Flavors and Fragrances and Givaudon. Stabilizers include
hydrated salts, such as magnesium sulfate, and boric acid.

In some of the compositions herein, adjuvants include (and are especially preferred) a chelating agent or sequestrant, most preferably, an aminopolyposphonate. These chelating agents assist in maintaining the solution stability of the salt activators and active oxygen source in order to achieve optimum performance. In this manner, they are acting to chelate heavy metal ions, which cause catalyzed decomposition of the (believed) in situ formed peroxyimide acids, although this is a non-binding theory of their action and not limiting.

The chelating agent is selected from a number of known agents which are effective at chelating heavy metal ions. The chelating agent should be resistant to hydrolysis and rapid oxidation by oxidants. Preferably, it should have an acid dissociation constant (pKa) of about 1-9, indicating that it dissociates at low pH's to enhance binding to metal cations. Acceptable amounts of the (optional) chelating agent range from 0-1,000, more preferably 5-500, most preferably 10-100 ppm chelating agent, in the wash liquor. As a hard surface cleaner, however, it is preferred to add amounts of the chelating agent from 0 - 100,000, more preferably 5 - 50,000, and most preferably 10-10,000 ppm chelating agent.

The most preferred chelating agent is an aminopolyposphonate, which is commercially available under the trademark Dequest from Monsanto Company. Examples thereof are Dequest 2000, 2041 and 2060. (See also Bossu U.S. Patent 4,473,507, column 12, line 63 through column 13, line 22, incorporated herein by reference.) A polyphosphonate, such as Dequest 2010, is also suitable for use.

Other chelating agents, such as ethylenediaminetetraacetic acid (EDTA) and nitrilotriacetic acid (NTA) may also be suitable for use. Still other new, preferred chelating agents are new propylenediaminetetraacetates, such as Hampshire 1,3 PDTA, from W.R. Grace, and Chel DTPA 100#F, from Ciba Geigy A.G. Mixtures of the foregoing may be suitable.

Additional desirable adjuncts are enzymes (although it may be preferred to also include an enzyme stabilizer). Proteases are one especially preferred class of enzymes. They are preferably selected from alkaline proteases. The term "alkaline," refers to the pH at which the enzymes' activity is optimal. Alkaline proteases are available from a

wide variety of sources, and are typically produced from various microorganism (e.g., Bacillus subtilis). Typical examples of alkaline proteases include Maxatase and Maxacal from International BioSynthetics. Alcalase, Savinase, and Esperase, all available from Novo Nordisk A/S. See also Stanislawski et al., U.S. Patent 4,511,490, incorporated herein by reference. Further suitable enzymes are amylases, which are carbohydrate-hydrolyzing enzymes. It is also preferred to include mixtures of amylases and proteases. Suitable amylases include Rapidase, from Société Rapidase, Milezyme from Miles Laboratory, and Maxamyl from International BioSynthetics.

Still other suitable enzymes are cellulases, such as those described in Tai, U.S. Patent 4,479,881, Murata et al., U.S. Patent 4,443,355, Barbesgaard et al., U.S. Patent 4,435,307, and Ohya et al., U.S. Patent 3,983,082, incorporated herein by reference.

Yet other suitable enzymes are lipases, such as those described in Silver, U.S. Patent 3,950,277, Thom et al., U.S. Patent 4,707,291, and Wiersema et al., U.S. Patents 5,296,161 and 5,030,240, and Poulou et al., U.S. Patent 5,108,457, incorporated herein by reference.

The hydrolytic enzyme should be present in an amount of about 0.01-5%, more preferably about 0.01-3%, and most preferably about 0.1-2% by weight of the detergent. Mixtures of any of the foregoing hydrolases are desirable, especially protease/amylase blends.

Some of the adjuncts, such as fluorescent whitening agents, enzymes and pigments, are sensitive to oxidants, and thus, may need to be co-dispensed in a separate liquid formulation. On the other hand, there are encapsulation methods and other protective additives available for these sensitive materials, such as, for example, from Coyne et al., U.S. Patents 4,863,626, 5,093,021, and 5,225,102 and DeLeeuw et al., U.S. Patents 5,254,287 and 5,167,854, incorporated herein by reference.

In some of the embodiments of this invention, such as when the delivery execution is a multiple delivery of liquid formulations, there may be a need for a viscosity/phase modifier. Exemplary such materials include alkanolamines, especially triethanolamine, and a wide variety of polymers, including water soluble to water-miscible polymers, such as polyethylene glycol, polyvinyl alcohol, polyvinyl

14
acetate, polyacrylic acid, co-polymers of acrylic acid, co-polymers of methacrylic acid, and the salts thereof. Other polymers include starch, xanthan gum, gum arabic and other naturally occurring polymers. Nonaqueous systems, on the other hand, can be thickened with silicas, such as xerogels and fumed and precipitated silicas, such as Cab-O-Sil.

5 Anti-redeposition agents, such as carboxy methylcellulose, are potentially desirable. Foam boosters, such as appropriate anionic surfactants, may be appropriate for inclusion herein. Also, in the case of excess foaming resulting from the use of certain surfactants, anti-foaming agents, such as alkylated polysiloxanes, e.g. dimethylpolysiloxane, would be desirable.

10 In certain hard surface cleaners, it is desirable to incorporate a source of particulate matter to act as abrasives. Abrasives are desirable adjuncts for cleaning especially persistent stains. Abrasives can be selected from a wide variety of particulate materials, including, but not limited to, calcium carbonate, glass beads, polymer beads, perlite, silica sand and various other insoluble, inorganic particulate abrasives are also
15 possible, such as quartz, pumice, feldspar, tripoli and calcium phosphate. See Brodbeck et al., U.S. Patent 5,529,711, and Choy et al., U.S. Patents 5,554,321 and 5,470,499, all of which are incorporated herein by reference. Other types of abrasives include water soluble materials present in an amount such as to exceed their solubility in water, leaving a portion thereof undissolved. These types of materials include alkali metal bicarbonates,
20 alkali metal phosphates, alkali metal borates, particularly sodium tetraborate decahydrate (borax) and pentahydrate (see U.S. Patent Application Serial No. 08/718,059, filed September 17, 1996, entitled "Cleaner with Water Soluble Abrasive," incorporated herein by reference.).

7. Liquid Medium

25 So that the correct amount of active ingredients are dosed, the inventive compounds are combined with a liquid medium, most preferably, an inert, nonreactive liquid. Generally speaking, this nonreactive liquid is principally water, or water combined with solvents, or nonaqueous solvents. For bench scale experiments, it is preferred to use deionized water, although in the large scale manufacture of the inventive
30 liquid compositions, this may not be necessary. The solvents can be chosen from any organic or inorganic solvents. Some preferred organic solvents are those with a vapor pressure of at least 0.001 mm Hg at 25°C and soluble to the extent of at least 1g/100ml

water. The organic solvents used in the invention are preferably selected from C_{1-6} alkanol, C_{3-24} alkylene glycol ether, and mixtures thereof. However, other, less water soluble or dispersible organic solvents may also be utilized. The alkanol can be selected from methanol, ethanol, n-propanol, isopropanol, butanol, pentanol, hexanol, their various positional isomers, and mixtures of the foregoing. It may also be possible to utilize in addition to, or in place of, said alkanols, the diols such as methylene, ethylene, propylene and butylene glycols, and mixtures thereof. Other solvents, such as amines, ketones, ethers, esters, carboxylic acids, oils, hydrocarbons and halides may be used alone, or mixtures thereof. In the case of certain amines, e.g., monoethanolamine, diethanolamine, etc., such solvents are also considered buffers (as described further above in 6). Thus, it is possible that, in certain instances, these amines can be bifunctional herein. Other examples of solvents can be found in Kirk-Othmer, Encyclopedia of Chemical Technology 3rd, Vol. 21, pp. 377-401 (1983), incorporated by reference herein. Where the formulation is a nonaqueous one, liquid nonionic surfactants, as previously mentioned, can be used to provide the continuous phase. Additionally, an inorganic or organic acid, such as listed above in 5, could be used as the continuous phase.

8. Applications

a. Laundry Products: Compositions of the invention are useful as or in laundry products, such as bleaching additives, detergents, detergent boosters, detergents with bleach, bleaches, bleaching aids, and stain removers. Among the advantages derived from compositions of the invention are improved cleaning, stain removal, spot removal, whitening, and brightening of treated articles.

Further benefits from use of the inventive compositions include scavenging of free dye during laundering to prevent dye transfer between garments (sometimes known as dye transfer inhibition).

b. Surface Cleaners: Other product applications include household cleaning products, such as hard surface cleaners to be dissolved in water prior to use. Exemplary surface cleaners are tile and grout cleaners, bathroom (floor, toilet, and counter) and kitchen (floor, sink, and counter) cleaners. Additionally, kitchen products such as dishwasher detergents with bleach or bleach cleaning and scrubbing pads are contemplated. Among the benefits derived from use of the inventive compositions in

16

such applications are improved stain and spot removal and general cleaning of the treated surfaces to remove food, rust, grime, mildew, mold, and other typical stains found on such surfaces.

5 In the hard surface cleaning area, the present invention has been found to benefit from more acidic conditions (i.e., lower pH's) than for laundry formulations. These pH's can be from about 0 to about 8, more preferably between about 2 and 7.5. At these lower pH levels, the extent of bleaching -- when the hard surface cleaner is formulated with both the MMA compound and a source of active oxygen -- appears to become improved.

10 In one preferred embodiment of a hard surface cleaner delivery, as in the laundry product formulation, a dual chambered container/dispenser is preferred. One chamber contained an H_2O_2 /MMA solution at an acidic pH. The other chamber contained agents to adjust the pH to the optimum level, namely, an alkaline buffer and optionally, a surfactant. Other agents could be included for improved cleaning performance (thickeners, chelating agents, builders, etc.) or aesthetic appeal (dyes, colorants, 15 fragrances). The preferred delivery is a trigger sprayer, which must blend the two solutions/fluids from the two chambers prior to delivering the combined solutions as a spray. This could be accomplished by either a mixing chamber, or by directing two fluid streams to a common target point by means of a diverter or other redirection means. The surfactant or surfactants used in the other chamber could be selected from the diverse variety disclosed above in 4. Further, it is common to include in hard surface cleaners at 20 least one solvent to further enhance cleaning performance and to disperse hydrophobic or poorly soluble materials into the liquid cleaner. Fragrances and colorants as provided from 6 are also desirable. Buffers selected from the various acids/alkaline materials disclosed from 5 are desirable, as well as thickening agents which allow the blended liquids to cling to vertical surfaces, such as colloids (clays, alumina, silica) or surfactants, 25 surfactant/solvent mixtures, or polymers. The thickening effect may be created in situ as the two liquid streams are blended at the nozzle or other delivery orifice, or pre-exist in thickened form. The molar ratio of active oxygen : MMA differs from its concentration in laundry product formulations in that a higher total amount is often present. For 30 example, the molar ratio can range from about 100:1 to about 1:10, more preferably about 10:1 to about 1:1, most preferably about 5:1 to about 1:1 H_2O_2 : MMA.

17

Additionally, non-household product applications are contemplated where an effective level of active oxygen generated in situ to treat water is useful. Illustrative of such applications are pool and spa additives, as well as cleaners to remove stains on outdoor concrete, stucco, siding, wood and plastic surfaces.

5 The advantages of a liquid cleaning or laundry product are quite evident. First, the MMA compounds are stable in hydrogen peroxide, allowing versatility in formulation. A single container or dual or multiple container delivery can be formulated. Liquid cleaning and laundering products appear to be greatly desired by consumers and it is advisable for the product formulator to offer liquid products as an alternative or preferred consumer
10 product. In laundry products, the liquid product is desirable to prespot or target stains on soiled fabrics and garments.

Aspects of the invention will now be illustrated by the following examples. It will be understood that these examples are intended to illustrate, and not to limit, the invention.

15

EXPERIMENTAL

EXAMPLE 1

20 In general, N-quaternary acetonitrile compounds are readily prepared from N-acetonitrile precursors by employing selected alkyl halides and using well-known synthetic approaches, such as are described by Menschutkin, *Z. Physik. Chem.*, 5, 589 (1890), and *Z. Physik. Chem.*, 6, 41 (1890); Abraham, *Progr. Phys. Org. Chem.*, 11, 1 (1974); and Arnett, *J. Am. Chem. Soc.*, 102, 5892 (1980). Further methods are depicted in German patent application DE OS 44 312 12, published March 7, 1996, and in the concurrently filed and co-pending application Serial No. _____, of inventors James E. Deline et al., entitled "Process for Preparing N-Alkyl Ammonium Acetonitrile
25 Compounds" (JSS Docket No.: 0409.073US___) both of which are incorporated herein by reference.

Compounds having the Formula I structure have a saturated ring formed by a plurality of atoms, broadly ranging from 3 to 9, although preferably containing 6 atoms

18

including the N₁ atom. Preparation of these compounds will most conveniently start with a compound already having the formed ring. For example, a number of preparations of inventive nitriles hereinafter described will begin with a morpholine (see, e.g., the Formula II structure). An example of three membered rings is aziridine, e.g.,

5 N-methylacetonitrile aziridinium; as an example of four membered rings there is azetidine, e.g., N-ethyl-N-methylacetonitrile azetidinium; as an example of five membered rings there is pyrrolidine, e.g., N-butylacetonitrile pyrrolidinium; as an example of six membered rings, in addition to morpholine, there is piperidine, e.g., acetonitrile piperidinium; as an example of seven membered rings there is

10 homopiperidinium; as an example of eight membered rings there is tropane, e.g., N-methylacetonitrile-8-azabicyclo[3.2.1]octane; and, as an example of nine membered rings there is octahydroindole, e.g., N-methylacetonitrile octahydroindolinium.

More particularly, in the preferred method of preparation a suitable amine is reacted with a monoaldehyde or a dialdehyde in an aqueous medium (Step A) followed by

15 subsequent quaternization (Step B) with an alkylating agent. In Step A, the reaction is preferably either in the pH range from 8 through 14, or the pH value is maintained at not less than 2 in Step B.

Thus, an amine with the formula



20 is reacted as Step A with a monoaldehyde or a dialdehyde R⁶-CHO or OHC-R⁵-CHO, whereby R⁵ is a chemical bond or a C₁ to C₆ alkylene bridge, or an oxyalkylene bridge, and R⁶ stands for H or C₁₋₂₀ alkyl, and with hydrogen cyanide or an alkali metal cyanide in an aqueous medium. Step B is quaternization with an alkylating agent R¹-X in an aqueous medium without isolating the intermediate product from Step A.

25 In Step A, cyanohydrins, e.g., formadehyde cyanohydrin, can be formed as by-products from the aldehyde, that is used, and hydrogen cyanide. These cyanohydrins do not react further with the alkylating agent in Step B so that renewed breaking down of the cyanohydrins into aldehyde and hydrogen cyanide in the final product is possible.

19

Without the procedure in accordance with the invention, Step B usually proceeds in a such a way that, as a result of hydrolysis of the added alkylating agent the pH value of the reaction mixture drifts off from the alkaline or neutral region into the strongly acidic region with increasing reaction time. The protonation of the amine nitrogen atom of the glycinonitrile, that has not yet been quaternized, sets in -- in competition with the alkylation -- starting from a certain pH value so that, at the end of the addition of the alkylating agent, no further reaction of the glycinonitrile takes place. Non-quaternized glycinonitrile in the final product can also represent an undesired source of hydrogen cyanide.

Step A generates especially good results if a pH range of 9 through 13 or, especially, 10 through 12, is utilized. In this pH range, the cyanohydrin that is formed is present in an equilibrium with the aldehyde and the hydrogen cyanide so that the re-formed adducts can react to completion with the amine to give glycinonitrile.

If one also uses an excess of amine that amounts to 2 through 20 mole % or, especially about 3 through 10 mole % or, most particularly of all, about 4 through 7 mole %, based on the quantity of the hydrogen cyanide or alkali metal cyanide that is used, then one achieves even more extensive suppression of hydrogen cyanide and ancillary components, that liberate hydrogen cyanide, in the final product.

Step B generates especially good results if the pH values are not reduced below 2.5 and, especially, not below 3. An optimum pH range for the quaternization of Step B is 2.5 through 5 or, especially, 3 through 4.

Use is also made of an excess of alkylating agent that amounts to 10 to 40 mole % or, especially, 15 to 25 mole % based on the quantity of amine that is used in Step A, then one achieves still more extensive suppression of the hydrogen cyanide and the subsidiary components, that liberate hydrogen cyanide, in the final product.

Once the nitriles are prepared in quaternized form, formation of the preferred bisulfate or sulfate form preferably is by heating an alkyl sulfate form, in an acid aqueous

20

solution. For example, a suitable elevated temperature is about 40° C to about 150°C; more preferably about 70°C to about 110°C. The acid aqueous solution may have a pH in the range of about -1 to 6, more preferably from 0 to 3, with the heating being for a period of about 1 to 50 hours.

5 Aspects of the invention will now be illustrated by the following examples. It will be understood that these examples are intended to illustrate, and not to limit, the invention.

EXAMPLE 2

Concentrated Laundry Additive Formulations

10 TABLE I discloses preferred, more preferred and most preferred ranges for a concentrated laundry additive which would be within the scope of this invention:

21

TABLE I

Ingredient	Preferred Range	More Preferred Range	Most Preferred
Anionic Surfactant	0-20%	0-10%	10.0%
Nonionic Surfactant	0-20%	0-10%	8.0%
Nonionic Ethoxylated Surfactant	0-20%	0-10%	6.5%
Amphoteric Surfactant	0-20%	0-10%	1.5%
Hydrogen peroxide	1-10%	3-6%	6%
MMA Compound ¹	0.1-10%	1-8%	2%
Fluorescent Whitening Agent	0-2%	0-1%	0.5%
Builder	0-10%	0-5%	1%
Preservative	0-0.1%	0-0.1%	0.02%
Chelating Agent	0-5%	0-2%	2%
pH Adjusting Salt ²	0-5%	0-2%	1%
Viscosity Modifier	0-10%	0.1-5%	3%
Dye/Colorant	0-0.5%	0-0.5%	0-0.5%
Fragrance	0-0.5%	0-0.5%	0-0.5%
Water	0-95%	q.s. to 100%	q.s. to 100%

¹Inventive compound, preferably MMAMS, MMAS, MMABS or MMATS.

²Can be either acid or alkali, depending on pH of formulation resulting from other ingredients. In general, it is desirable to attain, for optimal chemical storage stability, a final product pH from -1-5, more preferably 0-2 and most preferably 0-1. The acids and bases described in 5. above are utilized.

In the remaining examples, the same footnotes apply and will not be repeated.

22

In the next Example, a dual delivery system as discussed in 3., above, is disclosed which would be within the invention.

EXAMPLE 3

Two Part Concentrated Laundry Additive (Dual Chamber Bottle)

5

Part A (1st Chamber)

Ingredient	Preferred Range	More Preferred Range	Most Preferred
Anionic Surfactant	0-30%	0-20%	10.0%
Nonionic Surfactant	0-30%	0-20%	10.0%
Hydrogen Peroxide	1-10%	3-6%	6%
MMA	0.1-10%	1-8%	2%
Preservative	0-0.1%	0-0.1%	0.02%
Chelating Agent	0-5%	0-2%	2%
pH Adjusting Salt	0-5%	0-2%	1%
Dye/Colorant	0-0.5%	0-0.5%	0-0.5%
Water	0-95%	q.s. to 100%	q.s. to 100%

Part B (2nd Chamber)

Ingredient	Preferred Range	More Preferred Range	Most Preferred
Nonionic Ethoxylated Surfactant	0-30%	0-20%	6.5%
Amphoteric Surfactant	0-30%	0-20%	1.5%
Enzymes	0-5%	0-2%	1%
Fluorescent Whitening Agent	0-2%	0-1%	0.5%
Builder	0-10%	0-5%	1%
pH Adjusting Salt	0-10%	0-5%	0.5%
Viscosity Modifier	0-10%	0-5%	3%
Dye/Colorant	0-0.5%	0-0.5%	0-0.5%
Fragrance	0-0.5%	0-0.5%	0-0.5%
Water	0-95%	q.s. to 100%	q.s. to 100%

In the above system, Part A and Part B would be stored in separate compartments or chambers to prevent intermixing (cross-contamination or premature reaction). At time of use, preferably equal proportions (or, in an alternative embodiment, in a fixed ratio) of Part A and Part B would be combined by the user (for example, by co-metering the two

10

23
liquid formulations) and poured into, for example, a washing apparatus. Ingredients would be placed into the appropriate Part A or Part B so as to achieve the optimum physical and chemical storage stability, and individual ingredient stability. Additionally, by separate storage in a multiple delivery system, oxidation-sensitive materials, such as alkaline proteases, could be stored in an alkaline environment, while the acid-stable ingredients, such as hydrogen peroxide and the MMA compounds, could be stored in an acidic environment. Finally, as the MMA compound/hydrogen peroxide combination works best in an alkaline environment, upon dispensing into the alkaline wash water, the best performance conditions would be achieved upon such dispensing.

In the next Example, a preferred concentrated laundry additive formulation is disclosed which is within the scope of the invention.

Example 4
Concentrated Laundry Additive

Ingredient	Wt.% Active	Commercial Material
Anionic Surfactant	8%	C ₁₂₋₁₅ Alkylbenzene Sulfonate
Nonionic Surfactant	5%	Neodol 25-7
Hydrogen Peroxide	5%	Cosmetic Grade Peroxide
MMA Compound	3.5%	MMAMS*
Fluorescent Whitening Agent	0.2%	Tinopal CBS-X
Builder	0.5%	Sodium Chloride
Preservative	0.02%	Butylated Hydroxytoluene
Chelating Agent	2%	Dequest 2010
pH Adjusting Agent	1%	Sulfuric Acid
Viscosity/Phase Modifier	1%	Triethanolamine
Dye	0-0.5%	Various Vendors
Fragrance	0-0.5%	" "
D.I. Water	q.s.	

*Not a presently commercially available material.

In the next Example, an especially preferred two part concentrated laundry additive is disclosed which is within the invention.

24
Example 5

Two Part Concentrated Laundry Additive

Part A

Ingredient	Wt. %	Commercial Material
Anionic Surfactant	10.0%	C ₁₂₋₁₅ Alkylbenzene Sulfonate
Nonionic Surfactant	8.0%	Neodol 25-12
Hydrogen Peroxide	6.0%	Cosmetic Grade
MMA	2.0%	MMAMS
Preservative	0.2%	Butylated Hydroxytoluene
Chelating Agent	2%	Dequest 2006
pH Adjusting Salt	1%	Phosphoric Acid
Dye/Colorant	0-0.5%	Various Vendors
Water	q.s. to 100%	

Part B

Ingredient	Wt. %	Commercial Material
Nonionic Ethoxylated Surfactant	6.5%	Neodol 25-7
Nonionic Surfactant	1.5%	Barlox 12
Enzymes	1%	Novo Protease
Fluorescent Whitening Agent	0.5%	Tinopal AMS
Builder	1.0%	Sodium Chloride
pH Adjusting Salt	0.5%	Sodium Hydroxide
Viscosity Modifier	3%	Polyethylene Glycol
Dye/Colorant	0-0.5%	Various Vendors
Fragrance	0-0.5%	Various Vendors
Water	q.s. to 100%	

5 In Examples 6-8 below, exemplary dual chamber hard surface cleaner formulations were created and tested for stain removal performance against a laboratory mildew stain (*Aspergillus niger*). The stain was applied to bisque tiles (unglazed ceramic tiles to essentially mimic bathroom grout). A proprietary Minolta colorimeter was used to determine cleaning performance. To calibrate the colorimeter, a clean tile was used, and

10 then the stained tile was read. Then, after the inventive cleaning composition was applied, the colorimeter read the tiles, and the readout would indicate % stain removal at various elapsed time intervals. In the Examples, the pH level was varied to compare performance at differing pH levels. Part A contained the MMA compound, hydrogen

25

peroxide and a buffer. Part B contained an alkaline material.

Examples 6-8

Ingredient	Example 6			Example 7			Example 8		
	A	B	Blend	A	B	Blend	A	B	Blend
H ₂ O ₂	4.04		2.02	4.04		2.02	4.04		2.02
MMAMS	10		5	10		5	10		5
H ₂ SO ₄	2.43		1.21	0.89		0.45	0.89		0.45
NaOH								1	0.5
Na ₂ CO ₃		10	5		10	5		10	5
H ₂ O	balance	balance	balance	balance	balance	balance	balance	balance	balance
pH	7.6			8.5			9.5		
Molar Ratio of H ₂ O ₂ : MMAMS	3			3			3		

The formulations of Examples 6-8 were tested for performance in stain removal as described above. It was found that, at lower pH's, the formulation -- at equal dosages of active, i.e., at a 3:1 molar ratio of H₂O₂ : MMAMS -- had better stain removal performance at lower pH. This is graphically depicted in Fig. 1, in which the Y axis is % stain removal, the X axis is time in seconds. The data depicted have been adjusted for 100% stain removal by focusing on the area where the inventive bleaching formulation was applied.


It is to be understood that while the invention has been described above in conjunction with preferred specific embodiments, the description and examples are intended to illustrate and not limit the scope of the invention, which is defined by the appended claims.

26

Claims

1. A liquid cleaning composition including a compound with the structure of Formula I

FORMULA I



wherein A is a saturated ring formed by five atoms in addition to the N₁ atom, the five saturated ring atoms being four carbon atoms and a heteroatom, the substituent R₁ bound to the N₁ atom of the Formula I structure including either (a) a C₁₋₂₄ alkyl or alkoxyalkyl where the alkoxy is C₂₋₄, (b) a C₄₋₂₄ cycloalkyl, (c) a C₇₋₂₄ alkylaryl, (d) a repeating or nonrepeating alkoxy or alkoxyalkyl alcohol, where the alkoxy unit is C₂₋₄, or (e) -CR₂R₃C≡N where R₂ and R₃ are each H, a C₁₋₂₄ alkyl, cycloalkyl, or alkylaryl, or a repeating or nonrepeating alkoxy or alkoxyalkyl alcohol where the alkoxy unit is C₂₋₄, the R₂ and R₃ substituents are each H, a C₁₋₂₄ alkyl, cycloalkyl, or alkylaryl, or a repeating or nonrepeating alkoxy or alkoxyalkyl alcohol where the alkoxy unit is C₂₋₄, and wherein Y is an anionic counterion and Z is in the range of 0 to 10.

2. The liquid composition of claim 1 wherein said liquid includes an inert, nonreactive liquid.

3. The liquid composition of claim 2 wherein said liquid is selected from water, organic solvents and mixtures thereof.

4. The liquid composition of claim 1 wherein R₁ is a lower alkyl.

5. The liquid composition of claim 4 wherein R₁ is methyl.

6. The liquid composition of claim 5 wherein R₂ and R₃ are both hydrogen.

7. The liquid composition of claim 6 wherein the heteroatom is oxygen or sulphur.

8. The liquid composition of claim 7 wherein the heteroatom is oxygen.


27

9. The liquid composition of claim 1 wherein the counterion is either alkylsulfate, bisulfate, sulfate, tosylate or mixtures thereof.

10. The liquid composition of claim 1 wherein said liquid comprises a nonaqueous liquid.

5 11. The liquid composition of claim 10 wherein said nonaqueous liquid comprises liquid nonionic surfactants, organic solvents, or mixtures thereof.

12. A liquid bleaching composition, comprising:
a compound having the structure of Formula I



10 wherein A is a saturated ring formed by five atoms in addition to the N₁ atom, the five saturated ring atoms being four carbon atoms and a heteroatom, the substituent R₁ bound to the N₁ atom of the Formula I structure including either (a) a C₁₋₂₄ alkyl or alkoxyalkyl where the alkoxy is C₂₋₄, (b) a C₄₋₂₄ cycloalkyl, (c) a C₇₋₂₄ alkaryl, (d) a repeating or nonrepeating alkoxy or alkoxyalkyl alcohol, where the alkoxy unit is C₂₋₄, or
15 (e) -CR₂R₃C≡N where R₂ and R₃ are each H, a C₁₋₂₄ alkyl, cycloalkyl, or alkaryl, or a repeating or nonrepeating alkoxy or alkoxyalkyl alcohol where the alkoxy unit is C₂₋₄, the R₂ and R₃ substituents are each H, a C₁₋₂₄ alkyl, cycloalkyl, or alkaryl, or a repeating or nonrepeating alkoxy or alkoxyalkyl alcohol where the alkoxy unit is C₂₋₄, and wherein Y is an anionic counterion; and

20 a source of active oxygen.

13. The liquid bleaching composition as in claim 12 further including a chelating agent.

14. The liquid bleaching composition as in claim 12 further including one or more of a brightening agent and a whitening agent.

25 15. The liquid bleaching composition as in claim 12 further including at least

28

one enzyme.

16. The liquid bleaching composition as in claim 12 wherein R_1 of the Formula I compound is a lower alkyl.

17. The liquid bleaching composition as in claim 12 wherein the R_1 of the Formula I compound is methyl.

18. The liquid bleaching composition as in claim 12 wherein R_2 and R_3 of the Formula I compound are both hydrogen.

19. The liquid bleaching composition as in claim 18 wherein the heteroatom of the Formula I compound is oxygen.

20. The liquid bleaching composition as in claim 19 wherein the counterion of the Formula I compound is either alkylsulfate, bisulfate, sulfate, tosylate or mixtures thereof.

21. The liquid bleaching composition as in claim 12 wherein the source of active oxygen is selected from the group consisting of hydrogen peroxide, Caro's acid (peroxymonosulfuric acid), and then, as suspended particulate oxidants, the alkaline earth metal salts of percarbonate, perborate, persilicate, hydrogen peroxide adducts, and mixtures thereof.

22. The liquid bleaching composition as in claim 21 wherein the source of active oxygen is hydrogen peroxide.

23. The liquid bleaching composition as in claim 12 wherein said liquid bleaching composition is delivered in a dual chambered container.

24. The liquid bleaching composition as in claim 23 wherein the compound of Formula I and said source of active oxygen are delivered in a first chamber.

25. The liquid bleaching composition as in claim 23 wherein the compound of Formula I and said source of active oxygen are delivered in separate chambers.

26. The liquid bleaching composition as in claim 24 wherein first chamber contains a buffering material.

27. The liquid bleaching composition as in claim 26 wherein said buffering material comprises an acid.

5 28. The liquid bleaching composition as in claim 27 wherein the pH of the total composition varies from 0 to 10.

29. The liquid bleaching composition as in claim 24 wherein a second chamber contains an alkaline material.




FIG. 1

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/21335

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : 252/186.39; 510/312; 544/163, 86; 546/246

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 252/186.39; 510/312; 544/163, 86; 546/246

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONEElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)
APS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Hart, William F. et al. Some New Quaternary-Substituted Alkyl Morpholinium Chlorides and Pyrrolidinium alkyl Sulfates. The Journal Of Organic Chemistry. March 5, 1957, Vol. 22, No. 1, pages 86-88.	1-29
Y	US 4,978,770 A (AOYAGI ET AL) 18 December 1990 (18-12-90), see the abstract, column 7, lines 39-68 and examples 10-18.	1-29
Y	US 5,236,616 A (OAKES ET AL) 17 August 1993 (17-08-93), see the abstract, column 8, line 56 to column 10, line 10, and the examples.	1-29
A	US 4,397,757 A (BRIGHT ET AL) 09 August 1983 (09-08-83), see the abstract, and column 1, line 55 to column 2, line 18.	1-29

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

15 JANUARY 1998

Date of mailing of the international search report

05 MAR 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JOSEPH D. ANTHONY

Telephone No. (703) 308-0661

Form PCT/ISA/10 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/21335

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4,328,226 A (WITEK ET AL) 04 May 1982 (04-05-82), see the abstract, and the claims.	1-29

Form PCT/ISA/210 (continuation of second sheet)(July 1992)★

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/21335

A. CLASSIFICATION OF SUBJECT MATTER: IPC (6):

C01B 3/00; C11D 3/39, 7/38; C07D 265/12, 295/12, 295/14, 211/26, 211/34

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)